



Division of Child Care & Early Childhood Education  
P.O. Box 1437, Slot S150, Little Rock, AR 72203-1437  
P: 501.320.3971

---

## Notice of Incident

**Date of Incident:** 5/18/2022

**Date Reported to DCCECE:** 5/19/2022

**Agency Name:** Youth Home

**Agency Number:** 128

**Type of Facility:** PRTF

**Facility License Type:** Regular

**Type of Incident:** Emergency Room Visit

**Incident Description:** Client [REDACTED] reported to the nursing office at school c/o (R) arm pain where he has (49) stitches in (3) separate placements (upper arm around triceps brachia muscle and around lateral epicondyle area) after falling through a glass table prior to admission. Stitches around lateral epicondyle appear to begin to show wound dehiscence. MD notified with order to transport to ACH ER.

**Agency's Interim Corrective Action:** Transport to ER.

**Licensing Specialist Assigned:** C.DeBoer

**Licensing Supervisor Assigned:** 5/18/2022

---

**Child Abuse Hotline** (Only applies to maltreatment incidents)

**Was the Hotline Called:**No. **Was it accepted?**N/A **Outcome:** N/A

**Assigned Investigator:** N/A

---

**Date of DCCECE's Follow-up:** 5/19/2022 **Type of Follow-up:** Email

**Details from Follow-up:** Client [REDACTED] returned to campus from ACH ER. Results of exam: healing sutures. With instructions to wash daily, apply antibiotic ointment to wound. This will help

prevent thick scab from forming over sutures and keep covered. Sutures come out in 8 days from placement. Do not soak the wound. A small rim of pinkness or

redness around the edge of the wound is normal; however, it should not spread. Child may have mild tenderness or pain. Treat with Tylenol/ Motrin. Call your PCP or return to ED in 8-10 days

from 05/14/22 for suture removal. Seek medical attention sooner if any signs of infection such as redness, worsening pain, increased swelling, or drainage of pus.